

# Is there a role for retrobulbar hyaluronidase in hyaluronic acid vascular embolism related vision loss?

BY RACHNA MURTHY AND JONATHAN CP ROOS

## CPD record / Reader knowledge check

Please print out our questionnaire, complete the questions below and sign.

Keep for your records.

- Can hyaluronidase penetrate the optic nerve to dissolve HA embolus in the central retinal artery?
  - Yes
  - No
  
2. What visual assessments are recommended with HA filler treatments and vision loss? *(Tick all that apply)*
  - Pre-treatment visual acuity
  - Post vision loss visual acuity
  - Pupil reaction to light/RAPD
  - Visual field to confrontation
  
3. After diagnosis of HA related vision loss what measures can be beneficial for VISION RECOVERY? *(Tick all that apply)*
  - Ocular massage
  - Hypercapnoea
  - Reduce intraocular pressure
  - Retrobulbar hyaluronidase injection
  
4. Where should vision loss due to HA embolus be reported if possible? *(Tick all that apply)*
  - Ophthalmology emergency on call
  - Pharmaceutical company product specialist
  - MHRA
  - Register with [research@facorestoration.com](mailto:research@facorestoration.com) to be documented on global oculoplastic database

1 CPD POINT

Signature:

The CPD Certification  
 Service

Collective Mark

The **pmfa** Journal