

Temple filler: an unusual complication

BY RAED RTAIL, NICOLE DIB, KEVIN SAYEGH, RAYANE RAHI AND MICHELINE MAAMARI CHALOUHI

CPD record / Reader knowledge check

Please print out our form, complete the questions below and sign.

Keep for your records.

1. The largest vein coursing the temple and that is notorious to injury is:
 - ☐ Lateral temporal vein
 - ☐ Superficial temporal vein
 - ☐ Middle temporal vein
 - ☐ Ophthalmic vein
2. The drainage of the orbit depends mostly on:
 - ☐ The superior and inferior ophthalmic vein
 - ☐ Medial temporal vein
 - ☐ Lateral temporal vein
 - ☐ Supraorbital vein
3. Subconjunctival haemorrhage has been reported after:
 - ☐ Toxin injection to the corrugator
 - ☐ Filler injection in the cheek and tear trough
 - ☐ Periorbital toxin injection
 - ☐ Injury to the angular artery
4. The medial temporal vein is a small caliber vein:
 - ☐ True
 - ☐ False
5. When injecting the temple, the superficial temporal artery is the only risky vessel:
 - ☐ True
 - ☐ False

Signature:

CPD
CERTIFIED
The CPD Certification
Service
Collective Mark

1 CPD POINT

The **pmfa** Journal